

RESEARCH

Open Access



High prevalence of non-dipping patterns among Black Africans with uncontrolled hypertension: a secondary analysis of the CREOLE trial

Prossie Merab Ingabire^{1,2*}, Dike B. Ojji^{4,5}, Brian Rayner⁹, Elijah Ogola¹³, Albertino Damasceno¹⁴, Erika Jones⁹, Anastase Dzudie¹⁵, Okechukwu S. Ogah⁷, Neil Poulter¹¹, Mahmoud U. Sani⁸, Felix Ayub Barasa¹², Grace Shedul⁶, John Mukisa², David Mukunya^{2,3}, Bonnie Wandera², Charles Batte², James Kayima², Shahiemah Pandie¹⁰ and Charles Kiiza Mondo¹ CREOLE Study Investigators

Abstract

Background: Dipping of blood pressure (BP) at night is a normal physiological phenomenon. However, a non-dipping pattern is associated with hypertension mediated organ damage, secondary forms of hypertension and poorer long-term outcome. Identifying a non-dipping pattern may be useful in assessing risk, aiding the decision to investigate for secondary causes, initiating treatment, assisting decisions on choice and timing of antihypertensive therapy, and intensifying salt restriction.

Objectives: To estimate the prevalence and factors associated with non-dipping pattern and determine the effect of 6 months of three antihypertensive regimens on the dipping pattern among Black African hypertensive patients.

Methods: This was a secondary analysis of the CREOLE Study which was a randomized, single blind, three-group trial conducted in 10 sites in 6 Sub-Saharan African countries. The participants were 721 Black African patients, aged between 30 and 79 years, with uncontrolled hypertension and a baseline 24-h ambulatory blood pressure monitoring (ABPM). Dipping was calculated from the average day and average night systolic blood pressure measures.

Results: The prevalence of non-dipping pattern was 78% (564 of 721). Factors that were independently associated with non-dipping were: serum sodium > 140 mmol/l (OR = 1.72, 95% CI 1.17–2.51, *p*-value 0.005), a higher office systolic BP (OR = 1.03, 95% CI 1.01–1.05, *p*-value 0.003) and a lower office diastolic BP (OR = 0.97, 95% CI 0.95–0.99, *p*-value 0.03). Treatment allocation did not change dipping status at 6 months (McNemar's χ^2 0.71, *p*-value 0.40).

Conclusion: There was a high prevalence of non-dipping among Black Africans with uncontrolled hypertension. ABPM should be considered more routinely in Black Africans with uncontrolled hypertension, if resources permit, to help personalise therapy. Further research is needed to understand the mechanisms and causes of non-dipping pattern and if targeting night-time BP improves clinical outcomes.

Trial registration ClinicalTrials.gov (NCT02742467).

Keywords: Non-dipping pattern, Dipping pattern, Uncontrolled hypertension, Black African

*Correspondence: prossienkundiye@gmail.com

¹ St. Francis Hospital, Nsambya, Kampala, Uganda

Full list of author information is available at the end of the article



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

of Medicine, Faculty of Clinical Sciences, University of Abuja, Abuja, Nigeria. ⁵University of Abuja Teaching Hospital, Gwagwalada, Nigeria. ⁶Pharmacy, University of Abuja Teaching Hospital, Gwagwalada, Abuja, Nigeria. ⁷Cardiology Unit, Department of Medicine, University of Ibadan/University College Hospital, Ibadan, Nigeria. ⁸Department of Medicine, Bayero University, Aminu Kano Teaching Hospital, Kano, Nigeria. ⁹Division of Nephrology and Hypertension, Cape Town, South Africa. ¹⁰Hatter Institute of Cardiovascular Research in Africa, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa. ¹¹Imperial Clinical Trials Unit, School of Public Health, Imperial College London, London, UK. ¹²Department of Cardiology, Moi Teaching and Referral Hospital, Eldoret, Kenya. ¹³Department of Clinical Medicine and Therapeutics, University of Nairobi, Nairobi, Kenya. ¹⁴Eduardo Mondlane University Hospital, Maputo, Mozambique. ¹⁵Douala General Hospital, Douala, Cameroon.

Received: 11 March 2021 Accepted: 19 May 2021

Published online: 22 May 2021

References

- Collaborators GRF. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* (London, England). 2018;392(10159):1923.
- Adeloye D, Basquill C. Estimating the prevalence and awareness rates of hypertension in Africa: a systematic analysis. *PLoS ONE*. 2014;9(8):e104300.
- Ataklte F, Erqou S, Kaptoge S, Taye B, Echouffo-Tcheugui JB, Kengne AP. Burden of undiagnosed hypertension in sub-saharan Africa: a systematic review and meta-analysis. *Hypertension*. 2015;65(2):291–8.
- Pickering TG, Shimbo D, Haas D. Ambulatory blood-pressure monitoring. *N Engl J Med*. 2006;354(22):2368–74.
- O'Brien E, Parati G, Stergiou G, Asmar R, Beilin L, Bilo G, et al. European Society of Hypertension position paper on ambulatory blood pressure monitoring. *J Hypertens*. 2013;31(9):1731–68.
- Williams B, Mancia G, Spiering W, Agabiti Rosei E, Azizi M, Burnier M, et al. 2018 ESC/ESH Guidelines for the management of arterial hypertension: The Task Force for the management of arterial hypertension of the European Society of Cardiology and the European Society of Hypertension: The Task Force for the management of arterial hypertension of the European Society of Cardiology and the European Society of Hypertension. *J Hypertens*. 2018;36(10):1953–2041.
- Stergiou GS, Palatini P, Parati G, O'Brien E, Januszewicz A, Lurbe E, et al. 2021 European Society of Hypertension practice guidelines for office and out-of-office blood pressure measurement. *J Hypertens*. 2021;39:000.
- Turner JR, Viera AJ, Shimbo D. Ambulatory blood pressure monitoring in clinical practice: a review. *Am J Med*. 2015;128(1):14–20.
- Hermida RC, Smolensky MH, Ayala DE, Portaluppi F. Ambulatory Blood Pressure Monitoring (ABPM) as the reference standard for diagnosis of hypertension and assessment of vascular risk in adults. *Chronobiol Int*. 2015;32(10):1329–42.
- Angeli F, Reboldi G, Verdecchia P. Interpretation of ambulatory blood pressure profile: a prognostic approach for clinical practice. *J Hypertens*. 2015;33(3):454–7.
- Clement DL, De Buyzere ML, De Bacquer DA, de Leeuw PW, Duprez DA, Fagard RH, et al. Prognostic value of ambulatory blood-pressure recordings in patients with treated hypertension. *N Engl J Med*. 2003;348(24):2407–15.
- Mancia G, Verdecchia P. Clinical value of ambulatory blood pressure: evidence and limits. *Circ Res*. 2015;116(6):1034–45.
- Verdecchia P. Prognostic value of ambulatory blood pressure: current evidence and clinical implications. *Hypertension*. 2000;35(3):844–51.
- O'Brien E, Sheridan J, O'Malley K. Dippers and non-dippers. *Lancet*. 1988;2(8607):397.
- Lapinski M, Lewandowski J, Januszewicz A, Kuch-Wocial A, Symonides B, Wocial B, et al. Hormonal profile of dipper and non-dipper patients with essential hypertension. *J Hypertens Suppl Off J Int Soc Hypertens*. 1993;11(5):S294–5.
- Vaile JC, Stallard TJ, Al-Ani M, Jordan PJ, Townend JN, Littler WA. Sleep and blood pressure: spontaneous baroreflex sensitivity in dippers and non-dippers. *J Hypertens*. 1996;14(12):1427–32.
- Salles GF, Reboldi G, Fagard RH, Cardoso CR, Pierdomenico SD, Verdecchia P, et al. Prognostic effect of the nocturnal blood pressure fall in hypertensive patients: the ambulatory blood pressure collaboration in patients with hypertension (ABC-H) meta-analysis. *Hypertension*. 2016;67(4):693–700.
- Fagard RH, Thijs L, Staessen JA, Clement DL, De Buyzere ML, De Bacquer DA. Night-day blood pressure ratio and dipping pattern as predictors of death and cardiovascular events in hypertension. *J Hum Hypertens*. 2009;23(10):645–53.
- Cuspidi C, Sala C, Tadic M, Gherbesi E, De Giorgi A, Grassi G, et al. Clinical and prognostic significance of a reverse dipping pattern on ambulatory monitoring: an updated review. *J Clin Hypertens*. 2017;19(7):713–21.
- Timio M, Venanzi S, Lolli S, Lippi G, Verdura C, Monarca C, et al. "Non-dipper" hypertensive patients and progressive renal insufficiency: a 3-year longitudinal study. *Clin Nephrol*. 1995;43(6):382–7.
- Mahabala C, Kamath P, Bhaskaran U, Pai ND, Pai AU. Antihypertensive therapy: nocturnal dippers and nondippers. Do we treat them differently? *Vasc Health Risk Manag*. 2013;9:125–33.
- Wamba AA, Takah NF, Johnman C. The impact of interventions for the primary prevention of hypertension in Sub-Saharan Africa: a systematic review and meta-analysis. *PLoS ONE*. 2019;14(7):e0219623.
- Ojji DB, Mayosi B, Francis V, Badri M, Cornelius V, Smythe W, et al. Comparison of dual therapies for lowering blood pressure in Black Africans. *N Engl J Med*. 2019;380(25):2429–39.
- Ojji DB, Poulter N, Damasceno A, Sliwa K, Smythe W, Kramer N, et al. Rationale and design of the comparison of 3 combination therapies in lowering blood pressure in Black Africans (CREOLE study): 2 × 3 factorial randomized single-blind multicenter trial. *Am Heart J*. 2018;202:5–12.
- Mitu O, Roca M, Gurzu B, Magdalena M, Constantin L, Jitaru A, Al Namat R, Gavril RS, Mitu F. Predictors of the blood pressure non-dipping profile in newly diagnosed hypertensive patients. *J Hypertens Res*. 2016;2:121.
- Fagard R, Thijs L, Staessen JA, Clement D, De Buyzere M, De Bacquer D. Night-day blood pressure ratio and dipping pattern as predictors of death and cardiovascular events in hypertension. *J Hum Hypertens*. 2009;23(10):645–53.
- Tartan Z, Uyarel H, Kasikcioglu H, Alper AT, Ozay B, Bilsel T, et al. Metabolic syndrome as a predictor of non-dipping hypertension. *Tohoku J Exp Med*. 2006;210(1):57–66.
- Suzuki M, Guilleminault C, Otsuka K, Shiomi T. Blood pressure "dipping" and "non-dipping" in obstructive sleep apnea syndrome patients. *Sleep*. 1996;19(5):382–7.
- Jaeger BC, Booth JN III, Butler M, Edwards LJ, Lewis CE, Lloyd-Jones DM, et al. Development of predictive equations for nocturnal hypertension and nondipping systolic blood pressure. *J Am Heart Assoc*. 2020;9(2):e013696.
- Ben-Dov IZ, Kark JD, Ben-Ishay D, Mekler J, Ben-Arie L, Bursztyn M. Predictors of all-cause mortality in clinical ambulatory monitoring: unique aspects of blood pressure during sleep. *Hypertension*. 2007;49(6):1235–41.
- De La Sierra A, Redon J, Banegas JR, Segura J, Parati G, Gorostidi M, et al. Prevalence and factors associated with circadian blood pressure patterns in hypertensive patients. *Hypertension*. 2009;53(3):466–72.
- Ma Y, Sun S, Peng C-K, Fang Y, Thomas RJ. Ambulatory blood pressure monitoring in Chinese patients with obstructive sleep apnea. *J Clin Sleep Med*. 2017;13(3):433–9.
- Amer H, Awadi M, Shereef A, Mohammad M. Non-dipping phenomenon; Is it reversible in diabetic hypertensive cases. *J Clin Exp Cardiol*. 2017;8(565):2.
- Ayala DE, Moyá A, Crespo JJ, Castineira C, Dominguez-Sardina M, Gomara S, et al. Circadian pattern of ambulatory blood pressure in hypertensive patients with and without type 2 diabetes. *Chronobiol Int*. 2013;30(1–2):99–115.
- Fogari R, Zoppi A, Malamani GD, Lazzari P, Destro M, Corradi L. Ambulatory blood pressure monitoring in normotensive and hypertensive type 2 diabetics prevalence of impaired diurnal blood pressure patterns. *Am J Hypertens*. 1993;6(1):1–7.
- Hebert LA, Agarwal G, Ladson-Wofford SE, Reif M, Hiremath L, Carlton SG, et al. Nocturnal blood pressure in treated hypertensive African Americans

- compared to treated hypertensive European Americans. *J Am Soc Nephrol*. 1996;7(10):2130–4.
37. Muntner P, Lewis CE, Diaz KM, Carson AP, Kim Y, Calhoun D, et al. Racial differences in abnormal ambulatory blood pressure monitoring measures: results from the Coronary Artery Risk Development in Young Adults (CARDIA) study. *Am J Hypertens*. 2015;28(5):640–8.
 38. Mitu O, Roca M, Gurzu B, Magdalena M, Constantin L, Jitaru A, et al. Predictors of the blood pressure non-dipping profile in newly diagnosed hypertensive patients. 2016.
 39. Bankir L, Bochud M, Maillard M, Bovet P, Gabriel A, Burnier M. Night-time blood pressure and nocturnal dipping are associated with daytime urinary sodium excretion in African subjects. *Hypertension*. 2008;51(4):891–8.
 40. Sachdeva A, Weder AB. Nocturnal sodium excretion, blood pressure dipping, and sodium sensitivity. *Hypertension*. 2006;48(4):527–33.
 41. Fray J. Hypertension in blacks: physiological, psychosocial, theoretical, and therapeutic challenges. In: *Pathophysiology of hypertension in Blacks*. Springer; 1993. p. 3–22.
 42. Ergul A. Hypertension in black patients: an emerging role of the endothelin system in salt-sensitive hypertension. *Hypertension*. 2000;36(1):62–7.
 43. Rayner BL, Spence JD. Hypertension in blacks: insights from Africa. *LWW*; 2017.
 44. Ivy JR, Bailey MA. Nondipping blood pressure: predictive or reactive failure of renal sodium handling? *Physiology*. 2021;36(1):21–34.
 45. Burnier M, Coltai L, Maillard M, Bochud M, editors. *Renal sodium handling and nighttime blood pressure*. In: *Seminars in nephrology*. Elsevier; 2007.
 46. Osanai T, Okuguchi T, Kamada T, Fujiwara N, Kosugi T, Saitoh G, et al. Salt-induced exacerbation of morning surge in blood pressure in patients with essential hypertension. *J Hum Hypertens*. 2000;14(1):57–64.
 47. Uzu T, Ishikawa K, Fujii T, Nakamura S, Inenaga T, Kimura G. Sodium restriction shifts circadian rhythm of blood pressure from nondipper to dipper in essential hypertension. *Circulation*. 1997;96(6):1859–62.
 48. Wilson DK, Sica DA, Miller SB. Effects of potassium on blood pressure in salt-sensitive and salt-resistant black adolescents. *Hypertension*. 1999;34(2):181–6.
 49. Gregoire JR, editor. *Adjustment of the osmostat in primary aldosteronism*. In: *Mayo clinic proceedings*. Elsevier; 1994.
 50. Uzu T, Nishimura M, Fujii T, Takeji M, Kuroda S, Nakamura S, et al. Changes in the circadian rhythm of blood pressure in primary aldosteronism in response to dietary sodium restriction and adrenalectomy. *J Hypertens*. 1998;16(12):1745–8.
 51. Kusunoki H, Iwashima Y, Kawano Y, Hayashi S, Kishida M, Horio T, et al. Circadian hemodynamic characteristics in hypertensive patients with primary aldosteronism. *J Hypertens*. 2018;36(11):2260–8.
 52. Bochud M, Bovet P, Vollenweider P, Maillard M, Paccaud F, Wandeler G, et al. Association between white-coat effect and blunted dipping of nocturnal blood pressure. *Am J Hypertens*. 2009;22(10):1054–61.
 53. Zain-El Abidin M, Snincak M, Pahuli K, Solarova Z, Hrabcakova P. Non-dipping morning blood pressure and isolated systolic hypertension in elderly. *Bratisl Med J Bratisl Lek Listy*. 2013;114(3):150–4.
 54. Ben-Dov IZ, Perk G, Ben-Arie L, Mekler J, Bursztyn M. Pulse pressure is more susceptible to the white coat effect than is systolic blood pressure: observations from real-life ambulatory blood pressure monitoring. *Am J Hypertens*. 2004;17(6):535–9.
 55. Vasunta R-L, Kesäniemi YA, Ylitalo A, Ukkola O. Nondipping pattern and carotid atherosclerosis in a middle-aged population: OPERA Study. *Am J Hypertens*. 2012;25(1):60–6.
 56. Dai S, Huang B, Zou Y, Liu Y. Associations of dipping and non-dipping hypertension with cardiovascular diseases in patients with dyslipidemia. *Arch Med Sci AMS*. 2019;15(2):337.
 57. Hermida RC, Chayán L, Ayala DE, Mojón A, Fontao MJ, Fernández JR. Relationship between metabolic syndrome, circadian treatment time, and blood pressure non-dipping profile in essential hypertension. *Chronobiol Int*. 2011;28(6):509–19.
 58. Rorie DA, Rogers A, Mackenzie IS, Ford I, Webb DJ, Williams B, et al. Methods of a large prospective, randomised, open-label, blinded endpoint study comparing morning versus evening dosing in hypertensive patients: the Treatment In Morning versus Evening (TIME) study. *BMJ Open*. 2016;6(2):e010313.
 59. Hermida RC, Crespo JJ, Domínguez-Sardiña M, Otero A, Moyá A, Ríos MT, et al. Bedtime hypertension treatment improves cardiovascular risk reduction: the Hygia Chronotherapy Trial. *Eur Heart J*. 2020;41(48):4565–76.
 60. Kreutz R, Kjeldsen SE, Burnier M, Narkiewicz K, Oparil S, Mancia G. Blood pressure medication should not be routinely dosed at bedtime. We must disregard the data from the HYGIA project. *Blood Press*. 2020;29(3):135–6.
 61. Guthrie G, Poulter N, Macdonald T, Ford I, Mackenzie I, Findlay E, et al. Chronotherapy in hypertension: the devil is in the details. *Eur Heart J*. 2020;41(16):1606–7.
 62. Omboni S, Parati G, Palatini P, Vanasia A, Muiesan ML, Cuspidi C, et al. Reproducibility and clinical value of nocturnal hypotension: prospective evidence from the SAMPLE study. *J Hypertens*. 1998;16(6):733–8.
 63. Parati G, Staessen JA. Day–night blood pressure variations: mechanisms, reproducibility and clinical relevance. *J Hypertens*. 2007;25(12):2377–80.
 64. Ben-Dov IZ, Bursztyn M. Daytime sleeping and night-time urinating obscure normal dipping. *Nephrol Dial Transplant*. 2006;21(1):226–7.
 65. Perk G, Mekler J, Ishay DB, Bursztyn M. Non-dipping in diabetic patients: insights from the siesta. *J Hum Hypertens*. 2002;16(6):435–8.
 66. Stergiou G, Malakos J, Zourbaki A, Achimastos A, Mountokalakis T. Blood pressure during siesta: effect on 24-h ambulatory blood pressure profiles analysis. *J Hum Hypertens*. 1997;11(2):125–31.
 67. Bursztyn M, Mekler J, Wachtel N, Ben-Ishay D. Siesta and ambulatory blood pressure monitoring: Comparability of the afternoon nap and night sleep. *Am J Hypertens*. 1994;7(3):217–21.
 68. Matsumoto T, Tabara Y, Murase K, Setoh K, Kawaguchi T, Nagashima S, et al. Nocturia and increase in nocturnal blood pressure: the Nagahama study. *J Hypertens*. 2018;36(11):2185–92.
 69. Agarwal R, Light RP, Bills JE, Hummel LA. Nocturia, nocturnal activity, and nondipping. *Hypertension*. 2009;54(3):646–51.
 70. Leary AC, Donnan PT, MacDonald TM, Murphy MB. Physical activity level is an independent predictor of the diurnal variation in blood pressure. *J Hypertens*. 2000;18(4):405–10.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

